

WHAT IS CLAIMED IS:

1. A sustained release oral solid dosage form comprising:
 an therapeutically effective amount of a medicament having a solubility of more than about 10 g/l;
 a pH modifying agent;
 a sustained release matrix comprising a gelling agent, said gelling agent comprising a heteropolysaccharide gum and a homopolysaccharide gum capable of cross-linking said heteropolysaccharide gum when exposed to an environmental fluid, said dosage form providing a sustained release of said medicament after oral administration to human patients.
2. The sustained release oral solid dosage form of claim 1 which provides a sustained release of said medicament for at least about 12 hours after oral administration.
3. The sustained release oral solid dosage form of claim 1 which provides a sustained release of said medicament for at least about 24 hours after oral administration.
4. The sustained release oral solid dosage form of claim 1, wherein said medicament has a solubility of more than about 100 g/l.
5. The sustained release oral solid dosage form of claim 1, wherein said medicament has a solubility of more than about 1000 g/l.
6. The sustained release oral solid dosage form of claim 1, further comprising an inert pharmaceutical diluent.
7. The sustained release oral solid dosage form of claim 6, wherein said inert diluent is selected from the group consisting of monosaccharide, a disaccharide, a polyhydric alcohol, and mixtures thereof.
8. The sustained release oral solid dosage form of claim 6, wherein the ratio of said inert



ling agent
ned relea
d gelling
ned relea
gth enha
strength
olid dosa
n alkali r
lactate.
olid dosa
n sulfate.
olid dosa
n gum an

9. The sustained release oral solid dosage form of claim 1, wherein the ratio of said medicament to said gelling agent is from about 1:5 to about 5:1.

10. The sustained release oral solid dosage form of claim 1, further comprising an ionizable gel strength enhancing agent capable of crosslinking with said gelling agent and increasing the gel strength when the dosage form is exposed to an environmental fluid.

11. The oral solid dosage form of claim 10, wherein said ionizable gel strength enhancing agent comprises an alkali metal or an alkaline earth metal sulfate, chloride, borate, bromide, citrate, acetate, or lactate.

12. The oral solid dosage form of claim 11, wherein said ionizable gel strength enhancing comprises calcium sulfate.

13. The oral solid dosage form of claim 1, wherein said heteropolysaccharide gum comprises xanthan gum and said homopolysaccharide gum comprises locust bean gum.

14. ~~The oral solid dosage form of claim 1, wherein said pH modifying agent is an organic acid.~~

5. The sustained release oral solid dosage form of claim 14, wherein said organic acid is selected from the group consisting of citric acid, succinic acid, fumaric acid, malic acid, maleic acid, glutaric acid, lactic acid and combinations thereof.

16. The sustained release oral solid dosage form of claim 15, wherein said organic acid is fumaric acid.

17. The oral solid dosage form of claim 1, wherein said pH modifying agent is present in an amount from about 1% to about 10%.

540.1040

18. The oral solid dosage form of claim 1, further comprising a surfactant.
19. The oral solid dosage form of claim 18, wherein said surfactant is selected from the group consisting of anionic surfactants, cationic surfactants, amphoteric (amphipathic/amphiphilic) surfactants, and non-ionic surfactants.
20. The oral solid dosage form of claim 18, wherein said surfactant is selected from the group consisting of sodium lauryl sulfate and a pharmaceutically effective salt of docusate.
21. The oral solid dosage form of claim 1, wherein said sustained release matrix further comprises a hydrophobic material.
22. The oral solid dosage form of claim 21, wherein said hydrophobic material is selected from the group consisting of an alkylcellulose, a copolymer of acrylic and methacrylic acid esters, waxes, shellac, zein, hydrogenated vegetable oil, and mixtures thereof, in an amount effective to slow the hydration of said gelling agent when exposed to an environmental fluid.
23. The oral solid dosage form of claim 21, wherein said hydrophobic material is ethylcellulose.
24. The oral solid dosage form of claim 5, wherein said sustained release matrix comprises from about 1 to about 20% by weight of said hydrophobic material.
25. The oral solid dosage form of claim 1, further comprising from about 1 to about 10% by weight microcrystalline cellulose.
26. The oral solid dosage form of claim 1, wherein said medicament is a benzothiazine.
27. The oral solid dosage form of claim 26, wherein said benzothiazine is diltiazem or a pharmaceutically effective salt thereof.

540.1040

28. The oral solid dosage form of claim 27, which provides a sustained release of said diltiazem for at least about 12 hours after oral administration to human patients.
29. The oral solid dosage form of claim 28 wherein said diltiazem is present in an amount from about 60 mg to about 120 mg.
30. The oral solid dosage form of claim 27, which provides a sustained release of said diltiazem for at least about 24 hours after oral administration to human patients.
31. The oral solid dosage form of claim 30 wherein said diltiazem is present in an amount from about 120 mg to about 300 mg.
32. The oral solid dosage form of claim 1, wherein said medicament is an antispasmodic agent.
33. The oral solid dosage form of claim 32, wherein said antispasmodic drug is oxybutynin or a pharmaceutically acceptable salt thereof.
34. The oral solid dosage form of claim 33, wherein said antispasmodic agent is oxybutynin chloride.
35. The oral solid dosage form of claim 33, which provides a sustained release of said oxybutynin for at least 12 hours after oral administration to human patients.
36. The oral solid dosage form of claim 35, wherein said oxybutynin is present in an amount from about 2.5 mg to about 25 mg.
37. The oral solid dosage form of claim 33, which provides a sustained release of said oxybutynin for at least about 24 hours after oral administration to human patients.
38. The oral solid dosage form of claim 37, wherein said oxybutynin is present in an

540.1040

amount from about 5 mg to about 50 mg.

39. The oral solid dosage form of claim 1 which is a tablet.

40. The oral solid dosage form of claim 1 which is in granular form.

41. The oral solid dosage form of claim 40, wherein a portion of said medicament is outside the granulation.

42. The oral solid dosage form of claim 40, wherein a sufficient amount of said granules to provide an effective dose of said medicament are disposed in a pharmaceutically acceptable capsule.

43. The oral solid dosage form of claim 39 wherein at least part of a surface of said tablet is coated with a hydrophobic material to a weight gain from about 1 to about 20 percent, by weight.

44. The oral solid dosage form of claim 43, wherein said hydrophobic material is selected from the group consisting of an alkylcellulose, a copolymer of acrylic and methacrylic acid, waxes, shellac, zein, hydrogenated vegetable oils, and mixtures of any of the foregoing.

45. The sustained release oral dosage form of claim 18 which provides bimodal absorption profile of said medicament.

46. A sustained release oral solid dosage form comprising:

- an effective amount of a calcium channel blocker to provide a therapeutic effect, said calcium channel blocker having a solubility greater than 10 g/L;
- a pH modifying agent;
- a pharmaceutically acceptable surfactant
- a sustained release excipient comprising a gelling agent, said gelling agent comprising a heteropolysaccharide gum and a homopolysaccharide gum capable of cross-

540.1040

linking said heteropolysaccharide gum when exposed to an environmental fluid, said dosage form providing bimodal absorption profile of said calcium channel blocker and providing a sustained release of said calcium channel blocker for at least about 12 hours after oral administration to human patients.

47. The oral solid dosage form of claim 46, wherein said calcium channel blocker is a benzothiazine.

48. The oral solid dosage form of claim 46, wherein said benzothiazine is diltiazem or a pharmaceutically acceptable salt thereof.

49. The oral solid dosage form of claim 48, wherein said dosage form provides an initial peak concentration (C_{max} #1) of said diltiazem in about 4 to about 10 hours after oral administration of the dosage form, followed by a second peak concentration (C_{max} #2) which occurs in about 10 to about 16 hours after oral administration of the dosage form, said dosage form providing a therapeutic effect for at least about 24 hours after oral administration to a human patient.

50. The dosage form of claim 49, wherein said time to first peak plasma concentration (T_{max} #1) of diltiazem occurs in about 6 to about 8 hours after oral administration of the dosage form to the patient.

51. The dosage form of claim 49, wherein the maximum plasma concentration of diltiazem at the first T_{max} (C_{max} #1) is from about 50 to about 100 ng/ml, per administration of a 240 mg dosage of diltiazem.

52. The dosage form of claim 46, wherein the second peak plasma concentration (C_{max} #2) occurs in about 12 to about 14 hours after oral administration of the dosage form to the patient (T_{max} #2).

53. The dosage form of claim 52, wherein the maximum plasma concentration of

540.1040

diltiazem at Cmax #2 is from about 60 to about 90 ng/ml, per 240 mg diltiazem.

54. The dosage form of claim 46, wherein the width of the plasma concentration curve at 50% of the height of Cmax #1, based on a trough taken as the Cmin between Cmax #1 and Cmax #2 is from about 0.5 to about 4.0 hours.

55. The dosage form of claim 46, wherein the width of the plasma concentration curve at 50% of the height of Cmax #1, based on a trough taken as the Cmin between Cmax #1 and Cmax #2 is from about 1 to about 3 hours.

56. The dosage form of claim 46, wherein the width of the plasma concentration curve at 50% of the height of Cmax #2, based on a the trough taken as the Cmin between Cmax #1 and Cmax #2 is from about 0.5 to about 8 hours.

57. The dosage form of claim 46, wherein the width of the plasma concentration curve at 50% of the height of Cmax #2, based on a the trough taken as the Cmin between Cmax #1 and Cmax #2 is from about 2 to about 6 hours.

58. The dosage form of claim 49, wherein the ratio of Cmax #1 to Cmax #2 is from about 0.5:1 to about 1.5:1.

59. The dosage form of claim 58, wherein the ratio of Cmax #1 to Cmax #2 is from about 0.7:1 to about 1.2:1.

60. A method of treating a human patient suffering from a disease state selected from the group consisting of hypertension, angina, aneurysms, arhythmias, and headache comprising administering to said patient a dosage form of claim 46.

61. A sustained release oral solid dosage form comprising:
an effective amount of oxybutynin or a pharmaceutically acceptable salt thereof
to provide a therapeutic effect,

1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

a pH modifying agent;

a sustained release excipient comprising a gelling agent, said gelling agent

62. The sustained release oral solid dosage form of claim 61 which provides a time to peak plasma concentration (Tmax) in about 5 to about 15 hours

63. The sustained release oral solid dosage form of claim 62 which provides a time to peak plasma concentration (Tmax) in about 8 to about 12 hours.

64. A method of treating a human patient suffering from incontinence comprising administering to said patient a dosage form of claim 61.



Add Cl